

Variable Stars in the Sextans Dwarf Spheroidal Galaxy


... a progress report

Kathy Vivas

*Cerro Tololo Interamerican Observatory
La Serena, Chile*

Collaborators: M. Mateo (Michigan), J. Alonso-García (PUC, Chile), A. Walker (CTIO), D. Nidever (Michigan), B. Howard (CTIO REU student)


Dwarf Galaxies and their Variables




Updated map of MW satellites (DES Collaboration, 2015)

- Milky Way satellites allow studying the process of galaxy assembly
- SFH are different in many of these galaxies. Why?

The Role of Variable Stars




Carina dSph CMD (Monelli et al 2003)




- Anomalous Cepheids
- RR Lyrae Stars
- Dwarf Cepheids

The Importance of Dwarf Cepheids



Cohen et al (2012)

P-L relationship \rightarrow standard candles



*Amplitudes of dwarf cepheids in Carina,
Vivas & Mateo (2013)*


Only 1 DC (out of 1600) in Kepler field
has Amp > 0.1 mag

Observing Dwarf Cepheids




- Observational Challenges:
 - Faint
 - Short periods (~ 1 hour)
- Only 3 external galaxies have been searched for dwarf cepheids



The Sextans Dwarf Spheroidal Galaxy





- ~100 kpc
- Mostly old stars but also, intermediate age population
- Large galaxy, low surface brightness
- 2 DECam fields within the tidal radius
- Time series in g and r (~30 epochs/band, with a cadence of ~10 minutes)





Procedure

- Community Pipeline
- DoPhot photometry on individual images
- **Cleaning**
- Matching catalogs by coordinates
- Relative photometry
- Calibration
- Variables




Procedure

- Community Pipeline
- DoPhot photometry on individual images
- Cleaning
- Matching catalogs by coordinates
- Relative photometry
- Calibration
- Variables




Procedure

- Community Pipeline
- DoPhot photometry on individual images
- Cleaning
- Matching catalogs by coordinates
- Relative photometry
- Calibration
- **Variables**









Period Finding



Method based on Lafler and Kinman (1965) but using the 2-bands simultaneously



New Variables



Location of the variables in the CMD and in the sky



Summary and Future Work

- We have successfully put together a pipeline to identify short period variable stars
- So far, 33 RR Lyrae stars and 2 dwarf cepheids
- Sgr dwarf is next
- Study extension of the different stellar populations in these galaxies
- Deep CMDs over the whole area

